



Associates in Gastroenterology, P.C.
719-635-7321
www.agcosprings.com

Preparing for Your Procedure

Stop taking the following medications:

You will need to stop taking certain blood-thinning medications in the days before your colonoscopy. If you take a blood-thinning medication, please check with the doctor who prescribed your medication before stopping it.

7 Days Prior To Your Procedure:

- Stop taking all prescription anti-platelet medications: Plavix (clopidogrel), Pletal, Ticlid (ticlopidine), Aggrenox, Agrylin (anagrelide), or Effient.

5 Days Prior To Your Procedure:

- Stop taking Coumadin (warfarin), Brilinta, and Jantoven.

2 Days Prior To Your Procedure:

- Stop taking Pradaxa (dabigatran), Xarelto (rivaroxaban), Eliquis (apixaban), and oral iron supplements.

12 hours Prior To Your Procedure:

- Stop taking Lovenox, aspirin, ibuprofen, other NSAIDs.

Follow a special diet before the colonoscopy.

You will not be able to eat solid food for 2 days preceding your exam. Drinks are limited to **clear liquids** (water, sports drinks, tea, broth, jello, popsicles, clear fruit juices without pulp, coffee without milk or creamer, or clear carbonated beverages). Avoid red and purple liquids, which can be confused with blood during the colonoscopy.

Take a Bowel Prep.

Before a colonoscopy, you will need to clean out (empty) your colon. To accomplish this, you will need to follow the bowel preparation instruction on the other side of this page. Please follow the instructions provided on this form and disregard those on the prep box or given to you by the pharmacy. The laxative in your bowel preparation will cause you to have diarrhea. You should plan to be near a bathroom while taking your prep. Any residue in your colon may obscure the view of your colon during the exam.

Bring a medication list.

Tell your doctor about all the medications and supplements you are taking before your endoscopy. Be sure to bring a medication list with you. If you take heart, blood pressure, or seizure medication routinely, you may take the medication up to 2 hours before your procedure with a small amount of water. If you are diabetic, hold oral medications and short-acting insulin the morning of your procedure. Take only half of your normal dose of long-acting or basal insulin depending on your morning glucose level.

Plan ahead for your recovery.

You will receive anesthesia/sedation for this exam. You may feel mentally alert, but your memory, reaction times, and judgment may be impaired. You will not be allowed to drive or go back to work until the day after the procedure. You are not allowed to take a taxi, non-medical ride service, bus, or leave the Center alone. **IF YOU DO NOT HAVE A RESPONSIBLE ADULT TO ACCOMPANY YOU HOME, YOUR PROCEDURE WILL BE CANCELLED.**

Your Appointment is scheduled for:

Date: _____

Check-In Time: _____

Procedure Time: _____

Location:

Endoscopy Centers of Colorado Springs

- Audubon Medical Campus
2940 N. Circle Dr.
- St. Peregrine Pavilion
6031 E. Woodmen Rd. Ste. 100

Hospitals

- Penrose Main Hospital GI Lab
2222 N. Nevada Ave.
- St. Francis Hospital GI Lab
6001 E. Woodmen Rd.


If you are scheduled at a hospital:

- Call 776-8010 one (1) week prior to your procedure to pre-register
- Check in at registration –they will direct you to the GI Lab.

2 Day Miralax-Miralax: Bowel Preparation Instructions

Shopping List (these products can be purchased without a prescription):

- 1 (one) Bottle of Miralax powder (119 grams)
- 1 (one) Bottle of Miralax powder (238 grams)
- 3 (three) 28oz. bottles of electrolyte replacement solution such as Gatorade, Pedialyte, Powerade or Propel. Diabetics should choose a sugar free variety. Avoid red/purple coloring.



Two days before your colonoscopy:

Start a clear liquid diet.

No solid food starting in the morning. Begin a **clear liquid only** diet.

Between 4 PM and 6 PM:

Mix and drink Miralax – Gatorade preparation.

Step 1: Add the Miralax powder (119 grams) to 1 (one) 28 ounce bottle of electrolyte solution and mix/stir.

Step 2: Drink **ALL** 28 ounces of the solution by drinking 1 (one) 8 oz. glass approximately every 15 minutes until the mixture is gone.

Important: If you experience prep-related symptoms (for example, nausea, bloating, or cramping) pause or slow down the rate of drinking until your symptoms diminish.

Step 3: After completing the solution, you should continue with clear liquids of your choice to prevent dehydration.

One day before your colonoscopy:

At 6 PM on the day before your colonoscopy:

Mix and drink Miralax – Gatorade preparation.

Step 4: Add half of the 238-gram Miralax powder to 1 (one) 28 ounce bottle of electrolyte solution and mix/stir.

Step 5: Drink ALL 28 ounces of the solution over the next hour by drinking one glass approximately every 15 minutes.

Step 6: After completing the solution, you should continue with clear liquids of your choice between prep doses.

Eight hours prior to your colonoscopy:

Repeat steps 4 through 6. You must start your second dose at least (8) eight hours prior to your procedure time.

Continue to have clear liquids up to 6 hours prior to your procedure.

Six hours prior to your colonoscopy:

Stop drinking and have nothing more by mouth.

(This includes gum, lozenges, and hard candy.)

IF YOU DO NOT STOP DRINKING (6) SIX HOURS BEFORE YOUR PROCEDURE, YOUR PROCEDURE MAY BE CANCELLED.

FOR YOUR SAFETY, DO NOT SMOKE, VAPE, OR CHEW TOBACCO (6) SIX HOURS BEFORE YOUR PROCEDURE. DO NOT INJECT MARIJUANA THE DAY OF YOUR PROCEDURE.